

1 **Accepted for publication in *Freshwater Biology*, 19/01/2017**

2

3 **Trophic consequences for riverine cyprinid fishes of angler subsidies based on marine**  
4 **derived nutrients**

5

6 Catherine Gutmann Roberts<sup>1</sup>, Tea Bašić<sup>1</sup>, Fatima Amat Trigo<sup>1,2</sup>, J Robert Britton<sup>1\*</sup>

7

8 <sup>1</sup>Centre for Conservation Ecology and Environmental Sciences, Faculty of Science and  
9 Technology, Bournemouth University, Poole, Dorset, BH12 5BB, UK.

10 <sup>2</sup>Departamento de Zoología y Antropología Física, Universidad de Murcia, Spain

11

12

13

14

15

16

17 \*Corresponding author: [rbritton@bournemouth.ac.uk](mailto:rbritton@bournemouth.ac.uk). (+44)01202965384

18

## 19 Summary

20

- 21 1. The crossing of freshwater ecosystem boundaries by marine derived nutrients (MDN)  
22 is usually associated with migratory salmonid fishes returning to natal rivers. An  
23 alternative source of MDN in freshwaters is the widespread use of pelletized marine  
24 fishmeal ('pellets') by freshwater anglers as they target large bodied cyprinid fishes,  
25 such as European barbel *Barbus barbus*.
- 26 2. Here, the trophic consequences of MDN from pellets for riverine cyprinid fishes were  
27 tested. Approaches used stable isotope analyses in controlled and wild scenarios,  
28 using *B. barbus* and chub *Squalius cephalus* as model species. The isotopic niche,  
29 measured as standard ellipse area, was used to assess trophic niche size, and mixing  
30 models predicted the extent to which MDN contributed to fish diet.
- 31 3. In experimental mesocosms, *B. barbus* fed low volumes of pellets (approximately 3  
32 per fish) for 130 days had isotopic niche sizes that were up to four times larger than a  
33 control and 'medium' (6 per fish) and 'high' pellet (12 per fish) treatments. Somatic  
34 growth rates were significantly higher in the 'medium' and 'high' treatments. In pond  
35 enclosure experiments, when juvenile *B. barbus* and *S. cephalus* were fed pellets daily  
36 for 100 days, there was a substantial and significant shift in the position of their  
37 isotopic niche compared to controls with no pellets fed. However, for each species,  
38 there were no significant differences in their somatic growth rates in the presence/  
39 absence of pellets.
- 40 4. In a lowland river, high proportions of MDN contributed to the diet of *B. barbus* and  
41 *S. cephalus* captured by angling, but with substantial individual variability in those  
42 captured by electric fishing. Across all *B. barbus* > 400 mm, MDN dietary  
43 contributions ranged between 9 and 71%. This suggested some individual diet

44 specialisations within their population that was associated with feeding on this angler  
45 subsidy and that also resulted in a significant increase in the size of their population  
46 isotopic niche.

47 5. These results suggested that when pellets containing MDN are used in freshwater  
48 angling, they are consumed and assimilated by cyprinid fishes, influencing individual  
49 and population trophic positions, and isotopic niche sizes and dietary specialisations.  
50 The results also suggested that the extent to which individuals specialise in feeding on  
51 pellets potentially influences their vulnerability to capture by anglers.

52

53 **Keywords:** Allochthonous, barbel, fishmeal, MDN, river ecology, stable isotopes

54

55

56 **Introduction**

57

58 Trophic fluxes of energy and nutrient resources can be ecologically significant when they  
59 cross the boundaries of ecosystems that differ in their productivity (e.g. Polis & Hurd, 1995;  
60 Zhang *et al.*, 2003; Richardson *et al.*, 2016). These cross-system fluxes can maintain the  
61 productivity, diversity, and community structure of recipient ecosystems (Schindler *et al.*,  
62 2005). Anadromous salmonid fishes are well recognised as playing integral roles in these  
63 processes, as they accumulate the majority of their biomass in the ocean and import these into  
64 freshwaters during spawning, thus releasing marine derived nutrients (MDN) into the  
65 relatively nutrient-poor freshwater systems (Schindler *et al.*, 2003). However, this delivery  
66 mechanism is not the only MDN source in freshwaters, as aquaculture and angling activities  
67 can also elevate the quantity of MDN to freshwater ecosystems via the release of energy rich  
68 foods based on pelletized fishmeal ('pellets') that is derived from marine fishes (Bašić *et al.*,  
69 2015).

70

71 The use of marine derived fishmeal pellets in freshwater aquaculture is an integral part of the  
72 husbandry process (Naylor *et al.*, 2000). In recreational angling, marine derived fishmeal  
73 pellets of up to 21 mm in diameter are used as both an attractant and hook-bait, and thus they  
74 can supplement fish diet (Grey, Waldron & Hutchinson, 2004; Jackson *et al.*, 2013; Bašić *et*  
75 *al.*, 2015). These inputs of pellets can increase the productivity of freshwater systems due to  
76 their nutrient and energy fluxes (Jones *et al.*, 1998; Jefferies, 2000), and thus they can act as a  
77 strong allochthonous trophic subsidy (Marcarelli *et al.*, 2011; Sato & Watanabe, 2013). In  
78 doing so, they potentially alter food web structure via changes in the trophic interactions of  
79 consumers (Jefferies, 2000; Marzcak *et al.*, 2007), and potentially result in resource  
80 partitioning between populations (Bašić *et al.*, 2015). The pellets utilised by anglers tend to

81 have high protein levels from fishmeal (typically 40 to 50%) and lipid levels from fish oil  
82 (typically 20%) (Naylor *et al.*, 2000; Bašić *et al.*, 2015). These pellets have been used widely  
83 for at least 20 years by European freshwater anglers for exploiting the cyprinid fishes  
84 common carp *Cyprinus carpio* L. and European barbel *Barbus barbus* (L.) (Jackson *et al.*,  
85 2013; Bašić *et al.*, 2015). Substantial quantities can be used, with individual anglers often  
86 using in excess of 1 kg per day, with at least 10 anglers often being present daily on some  
87 small (< 1 km) stretches of English rivers in summer (Bašić *et al.*, 2015). Arlinghaus and  
88 Niesar (2005) estimated that the amount of bait used annually per freshwater angler in  
89 Germany was 7.3 kg, indicating that considerable volumes of angler bait might be introduced  
90 into freshwaters on an annual basis.

91

92 The provision of novel feeding opportunities, such as the seasonal availability of terrestrial  
93 insects for stream fishes (Syrjanen *et al.*, 2011), can result in individual trophic niche  
94 specialisation developing within populations (Britton & Andreou, 2016). This is where the  
95 population trophic niche consists of sub-groups of trophically specialised individuals that in  
96 entirety comprise the population niche (Araújo, Bolnick & Layman, 2011). The attractiveness  
97 of pelletized marine-derived fishmeal to many fishes is likely to relate to their provision of an  
98 energy rich resource that is relatively easy to assimilate and maximises growth rates (Naylor  
99 *et al.*, 2000; Bašić *et al.*, 2015). It was recently established that in four rivers in England, the  
100 diet of adult *B. barbus* comprised considerable proportions of pelletized fishmeal (up to 80%;  
101 Bašić *et al.*, 2015). However, this study was all based on samples collected from uncontrolled  
102 field conditions, with no consideration of how it impacted the population trophic niche of the  
103 fish or their somatic growth rates. The aim of this study was thus to quantify how MDN in  
104 pelletized fishmeal from angling modifies the population trophic niches, influences individual  
105 dietary specialisation, and affects the growth rates of riverine fishes. Following Grey,

106 Waldron & Hutchinson (2004) and Bašić *et al.* (2015), who established that MDN from  
107 pellets results in fish isotopic data being distinct within freshwater food webs, objectives  
108 were to: (1) assess how MDN modifies the trophic niche size and somatic growth rates of  
109 allopatric and sympatric fishes in controlled conditions; and (2) quantify the contribution of  
110 MDN to the diet of wild fishes, and assess its role in driving individual trophic niche  
111 specialisation and modification of the population trophic niche. It was hypothesised that  
112 where available, MDN pellets contribute substantial proportions of the diet of river fishes,  
113 resulting in individuals specialising on this trophic subsidy and having faster somatic growth  
114 rates.

115

## 116 **Materials and methods**

117

### 118 *Model species, experimental designs and field study*

119 The model species were *B. barbatus* and its cyprinid trophic analogue chub *Squalius cephalus*  
120 (L.). These fishes are sympatric in many European rivers and achieve relatively similar body  
121 sizes (Bašić & Britton, 2016). A mesocosm experiment tested how the variable availability of  
122 pellets affected the trophic niche size and somatic growth rates of allopatric *B. barbatus*. A  
123 semi-controlled pond experiment determined how pellet availability affected the trophic  
124 niche position and size, and somatic growth rates, of *B. barbatus* and *S. cephalus* in allopatry  
125 and sympatry. A field study then tested the influence of pellets on the trophic niche and diet  
126 composition of wild *B. barbatus* and *S. cephalus*. These studies utilised stable isotope analysis  
127 (SIA) to assess trophic niche sizes (as isotopic niches) and the diet composition of the fishes.

128

129 The mesocosm experiment was completed in 12 artificial ponds of 250 L volume, using  
130 hatchery-reared juvenile *B. barbatus* across four treatments: control (no supplementary

131 feeding), low (supplementary feeding of approximately three pellets per day per fish),  
132 medium (6 pellets per day per fish) and high (12 pellets per day per fish). Each treatment was  
133 replicated three times, with five fish used per replicate. The pellets were 2 mm diameter and  
134 constituent 45% protein (from marine fishmeal) and 20% fish oil (Dynamite Baits, 2017).  
135 Each mesocosm pond was outside, mounted on a concrete base with no overhanging trees  
136 nearby, and had a gravel substrate (6 mm diameter), aeration and a filter to maintain water  
137 quality. Feeding rates were achieved via automated feeders releasing pellets once per day at  
138 20:00, as *B. barbuis* are crepuscular (Britton & Pegg, 2011). The mesocosms were set up in  
139 April 2015 and were seeded with macroinvertebrates collected from a local stream  
140 (*Gammarus pulex*; 20 per mesocosm). Chironomid larvae naturally colonised all mesocosms.

141  
142 The fish were measured (fork length, nearest mm) and weighed (to 0.1 g) before their  
143 introduction into the mesocosms in June 2015 (Table 1). They were removed in October  
144 2015, thus were exposed to their new diets for 130 days. Temperature loggers (TinyTag TGP-  
145 4017) in eight mesocosms (2 per treatment) recorded water temperatures twice per day (0.00  
146 and 12.00) revealed a mean water temperature ( $\pm$  95% confidence limits) of  $19.4 \pm 0.7$  °C,  
147 with no significant differences between mesocosms (ANOVA:  $F_{1,6} = 0.56$ ,  $P = 0.48$ ). For a  
148 consumer of starting weight 10 g, estimated half-life at 20 °C is 36 days for  $\delta^{13}\text{C}$  and 38 days  
149 for  $\delta^{15}\text{N}$  (Thomas & Crowther, 2015). These values equate to 92% replacement of both  
150 isotopes in the fish after 130 days, with consumers generally considered to have fully  
151 equilibrated to their food resources at 94% isotopic replacement (Hobson & Clark, 1992).

152  
153 On day 130, the mesocosms were drained and the fish removed, euthanized (over-  
154 anaesthesia; MS-222), re-measured, re-weighed and a dorsal muscle sample taken for SIA  
155 (Busst, Bašić & Britton, 2015). Samples of putative prey resources were also collected from

156 each mesocosm (*G. pulex* and Chironomid larvae); where possible, these represented  
157 triplicate samples per mesocosm (1 sample = 5 individuals). All samples were then oven  
158 dried to constant weight at 60°C as preparation for SIA.

159

160 The pond experiment used mesocosms where *B. barbuis* and *S. cephalus* were used in  
161 allopatry and sympatry. Thus, three treatments were used in pellet presence and absence: both  
162 species in allopatry (n = 10), and a final treatment where they were present in sympatry (n = 5  
163 + 5), with three replicates per treatment. All fish were juveniles (starting lengths 60 to 88 mm,  
164 starting weights < 10 g) and hatchery reared. Each mesocosm was set up as per Bašić and  
165 Britton (2016), thus each comprised of an independent enclosure situated within one of two  
166 larger semi-natural, ex-aquaculture ponds (pond size: 30 x 12 m; consistent 1 m depth). Each  
167 enclosure comprised of aluminium frames of 1.66 m (length) x 1.05 m (width) x 1.2 m  
168 (height) within a net of 7 mm square mesh that prevented fish ingress/ egress but enabled  
169 transfer of water and invertebrates. The enclosures provided uniform habitats across the  
170 treatments and replicates in which the fish were exposed to the same prey communities. The  
171 enclosures in which pellets were fed were located in a separate pond to those with no pellets  
172 fed to avoid risk of cross-contamination between treatments. Within their larger ponds, the  
173 enclosures were located randomly, with least 0.5 m distance between them for independence.  
174 Water temperatures were measured hourly using a temperature logger (TinyTag TGP-4017)  
175 placed in the centre of each pond; mean temperature ( $\pm$  95% confidence limits) was  $18.2 \pm$   
176  $0.3$  °C in the non-pellet pond and  $18.4 \pm 0.4$  °C in the pellet pond. Anti-predator netting (15  
177 mm mesh) was also placed over the top of all enclosures. The enclosures sat on the substrate  
178 and macrophytes grew through each of them (primarily *Elodea* spp.)

179



180 The enclosures were placed into the ponds seven days before the fish were introduced, with  
181 the experimental period commencing in May 2014 and lasting 100 days. The estimated  
182 isotopic turnover was approximately 90% (Thomas & Crowther, 2015). Feeding of pellets  
183 used two methods. Firstly, 2 mm pellets were fed via automated feeders (30 per day).  
184 Secondly, 3 mm pellets were fed once per week by hand (approximately 60 pellets per  
185 replicate). Other than size, the pellets were identical to those used in the first mesocosm  
186 experiment, with the same ingredients and constituents (i.e. fishmeal-based, with the same  
187 protein and lipid levels; Dynamite Baits, 2017). Following the removal of the enclosures on  
188 day 100, the fish were recovered, euthanized (anaesthetic overdose, MS-222) and placed on  
189 ice, with samples of macroinvertebrates taken from each enclosure. In the laboratory, fish  
190 were re-measured and dorsal muscle samples taken. Macroinvertebrate samples were sorted  
191 to species, enabling three samples per species to be dried for SIA (Bašić & Britton, 2016). A  
192 random selection of fish dorsal muscle samples (n = 15 to 18 per species and treatment;  
193 minimum number of samples per replicate = 5) was then also selected and dried for SIA.

194

195 The field study used the invasive *B. barbuis* and native *S. cephalus* populations of the River  
196 Teme, Worcester (52°10'13" N; 2°14'31" W) to test the influence of MDN from pellets on the  
197 diet composition and trophic niche size of wild fishes. The study stretch receives considerable  
198 angling pressure for *B. barbuis* from both banks throughout the year, but especially between  
199 June and October when anglers are present daily, with the majority utilising pellets based on  
200 fishmeal. A previous study also indicated *B. barbuis* diet elsewhere on the river  
201 (approximately 10 km upstream, with separation by a weir of approximately 2.0 m head)  
202 consisted of high proportions of pelletized fishmeal (Bašić *et al.*, 2015). Here, SIA of the  
203 fishes utilised scales as only catch and release angling is practised for cyprinid fishes on the

204 river and so the collection of SIA material had to be rapid and non-destructive, but also  
205 appropriate for analysis (Hutchinson & Trueman, 2006; Busst & Britton, 2016).

206

207 Samples of *B. barbuis* were captured using a combination of boat mounted electric fishing on  
208 the 22<sup>nd</sup> September 2015 and angling on the 22<sup>nd</sup> and 23<sup>rd</sup> September. Samples of *S. cephalus*  
209 were captured by angling between 22<sup>nd</sup> and 30<sup>th</sup> September 2015. Fish were tagged with  
210 passive integrated transponder tags before their release, with no tagged fish recaptured. Each  
211 captured fish was measured (fork length ( $L_f$ ), nearest mm) and three to five scales removed  
212 and stored in paper envelopes. Concomitantly, samples of angler bait were taken for SIA.  
213 Samples of macroinvertebrates for SIA were collected by kick-sampling. This also provided  
214 samples of minnow *Phoxinus phoxinus*, bullhead *Cottus gobio* and stone loach *Barbatula*  
215 *barbatula* for SIA (hereafter referred to as ‘small fishes’; all were <40 mm). Triplicate  
216 samples were taken of each species, with dorsal muscle samples taken from each ‘small fish’.  
217 For SIA, the large body size (> 270 mm) of the sampled *B. barbuis* and *S. cephalus* meant that  
218 only material from the very outer portions of scales were used in analyses, i.e. material  
219 produced from recent growth (Hutchinson & Trueman, 2006; Bašić *et al.*, 2015).

220

#### 221 *Stable isotope analysis*

222 SIA of all samples was completed at the Cornell Isotope Laboratory, New York, USA, where  
223 the dried samples were ground to powder and weighed precisely to ~1000 µg in tin capsules  
224 and analysed on a Thermo Delta V isotope ratio mass spectrometer (Thermo Scientific, USA)  
225 interfaced to a NC2500 elemental analyser (CE Elantach Inc., USA). Verification for  
226 accuracy was against internationally known reference materials and calibrated against the  
227 primary reference scales for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ . Accuracy and precision of the sample runs was  
228 tested every 10 samples using a standard animal sample (mink). Overall standard deviation

229 was 0.11‰ for  $\delta^{15}\text{N}$  and 0.09 for  $\delta^{13}\text{C}$ , and analytical precision associated with the  $\delta^{15}\text{N}$  and  
230  $\delta^{13}\text{C}$  sample runs was estimated at 0.42 and 0.15‰ respectively. Data outputs were in delta  
231 ( $\delta$ ) isotope ratios expressed per mille (‰). No lipid correction was applied as C:N ratios  
232 indicated very low lipid content (Post *et al.*, 2007).

233

234 In the pond experiment, the 95% confidence limits of the mean SI data for the  
235 macroinvertebrates suggested some significant differences between the two larger ponds  
236 (‘pellet pond’:  $\delta^{13}\text{C}$ :  $-31.86 \pm 1.06$ ,  $\delta^{15}\text{N}$ :  $5.9 \pm 0.66$ ‰; ‘non-pellet pond’:  $\delta^{13}\text{C}$ :  $-34.68 \pm 1.14$ ,  
237  $\delta^{15}\text{N}$ :  $8.49 \pm 0.60$ ‰). Therefore, to enable true comparison between the pellet and no pellet  
238 treatments, the  $\delta^{15}\text{N}$  data were transformed to trophic position (TP), using the equation:

$$239 \text{TP}_i = [(\delta^{15}\text{N}_i - \delta^{15}\text{N}_{\text{base}})/3.4] + 2$$

240 where  $\text{TP}_i$  is the trophic position of the individual fish,  $\delta^{15}\text{N}_i$  is the isotopic ratio of that fish,  
241  $\delta^{15}\text{N}_{\text{base}}$  is the isotopic ratio of the primary consumers (macroinvertebrates), 3.4 is the  
242 fractionation between trophic levels and 2 is the trophic position of the baseline organism  
243 (Post, 2002). The  $\delta^{13}\text{C}$  data were converted to  $\delta^{13}\text{C}_{\text{corr}}$  using:

$$244 \delta^{13}\text{C}_{\text{corr}} = \delta^{13}\text{C}_i - \delta^{13}\text{C}_{\text{meaninv}}/\text{CR}_{\text{inv}}$$

245 where  $\delta^{13}\text{C}_{\text{corr}}$  is the corrected carbon isotope ratio of the individual fish,  $\delta^{13}\text{C}_i$  is the  
246 uncorrected isotope ratio of that fish,  $\delta^{13}\text{C}_{\text{meaninv}}$  is the mean invertebrate isotope ratio (the  
247 ‘baseline’ invertebrates) and  $\text{CR}_{\text{inv}}$  is the invertebrate carbon range ( $\delta^{13}\text{C}_{\text{max}} - \delta^{13}\text{C}_{\text{min}}$ ;  
248 Olsson *et al.*, 2009). As stable isotope data from dorsal muscle more closely reflects diet  
249 (Grey *et al.*, 2009), then for the fish samples from the field study, their SI scale data were  
250 converted to dorsal muscle tissue values before further analysis using conversion values from  
251 Busst, Bašić & Britton (2015) that are specific to *B. barbuis* and *S. cephalus*.

252

253 *Testing of stable isotope analysis data*

254 In all cases, the SI data were used to calculate the trophic niche sizes of the fishes, using the  
255 isotopic niche. The isotopic niche varies slightly from the trophic niche through factors  
256 including growth and metabolic rate of individuals, and thus is used here as an approximation  
257 of the trophic niche (Jackson *et al.*, 2011). It was measured using the metric ‘standard ellipse  
258 area’ (SEA), a bivariate measure of the distribution of individuals in trophic space (Jackson *et*  
259 *al.*, 2012). Each ellipse enclosed ~40% of the data and thus represents the typical resource  
260 use within the study population (Jackson *et al.*, 2011; Jackson *et al.*, 2012). Due to relatively  
261 small sample sizes, a Bayesian estimate of SEA (SEA<sub>b</sub>) was used that utilises a Markov chain  
262 Monte Carlo simulation with 10<sup>4</sup> iterations for each group and provides 95% confidence  
263 limits of isotopic niche size (Jackson *et al.*, 2011; R Core Team, 2014). Where appropriate, to  
264 indicate how similar fish isotopic niches were in MDN presence/ absence, the extent of niche  
265 overlap was also estimated (%).

266

267 Bayesian mixing models then estimated the relative proportions of different food resources  
268 contributing to fish diet using the MixSIAR package in R (Parnell *et al.*, 2010; R Core  
269 Development Team, 2013; Stock & Semmens, 2013). Correct for isotopic fractionation  
270 between resources and consumers used species-specific and tissue-specific fractionation  
271 factors between fish and prey ( $\delta^{15}\text{N}$ :  $3.4 \pm 0.98\text{‰}$ ;  $\delta^{13}\text{C}$ :  $0.39 \pm 1.3\text{‰}$ ) (Busst, Bašić &  
272 Britton, 2015; Busst & Britton, 2016). All models were run using normal run length (chain  
273 length: 100,000 iterations with burn-in of 50,000, with posterior thinning (thin: 50) and 3  
274 chains). Model diagnostics were based on Gelman-Rubin and Geweke, with sufficient  
275 convergence to accept the results (Stock & Semmens, 2013). In mesocosm experiments,  
276 models were run with the resources as ‘pellets’ and ‘macroinvertebrates’. The latter was  
277 primarily Chironomid larvae, as this was the only putative food resource sampled from each

278 individual mesocosm. However, it also covered *G. pulex*, as some samples were collected  
279 from a small proportion of the mesocosms. Their SI data overlapped with Chironomids and  
280 so the model could not separate their dietary contributions (mean SI values  $\pm$  95% confidence  
281 limits (%): Chironomid: n = 18;  $\delta^{13}\text{C}$ :  $-24.08 \pm 0.36$ ,  $\delta^{15}\text{N}$ :  $7.83 \pm 0.38$ ; *G. pulex*: n = 6;  $\delta^{13}\text{C}$ :  
282  $-23.78 \pm 0.46$ ,  $\delta^{15}\text{N}$ :  $8.29 \pm 0.24$ ). In the pond experiments, four putative food resources were  
283 used: 2 mm pellet, 3 mm pellet and the macroinvertebrate groups Corixidae and Odonata. In  
284 the field study, the putative food resources in the model were pooled according to fish pellet  
285 1, fish pellet 2, small fishes and Arthropoda (*cf.* Bašić *et al.*, 2015). In addition to the  
286 Bayesian mixing models already outlined, these field study data were then also used to assess  
287 individual variability using SOLOSIAR ('siarsolomcmc4') in the SIAR package in R  
288 (Parnell *et al.*, 2010; R Core Development Team, 2013). In this model, fractionation values  
289 were (mean  $\pm$  SD):  $\delta^{13}\text{C}$ :  $2.57 \pm 0.06$  for 'small fishes' and both pellets, and  $0.80 \pm 0.30$  for  
290 Arthropoda;  $\delta^{15}\text{N}$ :  $2.4 \pm 0.07$  for 'small fishes' and both pellets, and  $3.0 \pm 0.02$  for  
291 Arthropoda (Busst, Bašić & Britton, 2015; Busst & Britton, 2016).

292

### 293 *Other data analyses*

294 In the mesocosm and pond experiments, SI data were also tested in linear mixed effect  
295 models (LMEM). In the mesocosm experiment, differences were tested in the isotopic data of  
296 *B. barbus* between the four treatments. The dependent variable was  $\delta^{13}\text{C}$  or  $\delta^{15}\text{N}$ , and each  
297 model was fitted with mesocosm number as a random effect on the intercept to prevent  
298 inflation of the residual degrees of freedom (Tran *et al.*, 2015). The significance of  
299 differences in SI data between treatments used estimated marginal means and linearly  
300 independent pairwise comparisons with Bonferroni correction for multiple comparisons. In  
301 the pond experiment, differences were tested between the species, their allopatric and  
302 sympatric treatments, and between the pellet and no pellet treatments. Species were entered

303 into models according to their treatments so, for example, *B. barbuis* was present in models as  
304 (1) allopatric *B. barbuis*, (2) in sympatry with *S. cephalus*, and (3) in the presence and absence  
305 of pellets. The dependent variable was Ccorr or TP, with each model also fitted with  
306 mesocosm number as a random effect. The significance of differences in Ccorr and TP were  
307 also determined from the model outputs using linearly independent pairwise comparisons.

308

309 Somatic growth rates were estimated in the mesocosm experiments using incremental length  
310 (IL) and specific growth rate (SGR); IL was determined per replicate for each treatment and  
311 was expressed as the mean daily growth increment per fish. It was calculated from:

$$312 \quad [((\text{total } L_{t+1}) - (\text{total } L_t))/4]/t$$

313 where total  $L_t$  and  $L_{t+1}$  was the total starting and end lengths of the fish in each replicate, 4  
314 represents the number of fish per replicate and  $t$  = number of days. Mean specific growth  
315 rates (SGR) were determined from:

$$316 \quad 100[(\ln W_{t+1}) - (\ln W_t)]/4t$$

317 where  $W_t$  = total starting weight and  $W_{t+1}$  = total end weight. In the pond experiments, only  
318 incremental length was tested. Using generalised linear models, differences were tested in the  
319 growth rate of each species according to their context (allopatric or sympatric) and treatment  
320 (pellet or no pellet). In the field study, the scales of the fish were viewed on a projecting  
321 microscope and an age estimate derived. Scales measurements of total scale radius (SR) and  
322 distance to the penultimate and final annulus (PA and FA respectively) were then taken to  
323 enable the last annual length increment ( $L_{fa}$ ) of the fish to be calculated from:

$$324 \quad L_{fa} = ([FA-PA]/SR) \times L_f.$$

325 Throughout the results, where error is expressed around the mean, it represents 95%  
326 confidence limits unless stated otherwise.

327 **Results**

328

329 *Mesocosm experiments*

330 There were no significant differences in starting lengths and weights of the fish across the  
331 experimental treatments (generalized linear models: length: Wald  $\chi^2 = 0.91$ ,  $P = 0.47$ ; weight:  
332 Wald  $\chi^2 = 0.79$ ,  $P = 0.51$ ). At the conclusion of the experiment, all of the fish were  
333 recovered, and their mean length and weight had increased to  $120.4 \pm 4.1$  mm and  $18.3 \pm 2.0$   
334 g, with significant differences in final lengths and weights across the treatments (generalized  
335 linear model: Wald  $\chi^2 = 50.64$ ,  $P < 0.001$ ). Fish had higher lengths and mass in the Low,  
336 Medium and High treatments compared with the Control ( $P < 0.001$ ). The generalized linear  
337 model for both SGR and IL was significant (Wald  $\chi^2 = 263.9$ ,  $P < 0.001$  and Wald  $\chi^2 =$   
338  $2776.3$ ,  $P < 0.001$  respectively), with growth rates being significantly faster in all treatments  
339 compared with the Control ( $P < 0.001$ ; Fig. 1). Both SGR and IL increased as the proportion  
340 of pellets fed daily increased (Fig. 1).

341

342 The LMEM revealed significant differences in  $\delta^{13}\text{C}$  between *B. barbuis* in the control (mean -  
343  $21.4 \pm 0.17\text{‰}$ ) and the other treatments (Low:  $-21.7 \pm 0.2\text{‰}$ ; Medium:  $-22.1 \pm 0.1\text{‰}$ ; High: -  
344  $22.1 \pm 0.1\text{‰}$ ) ( $P < 0.001$ ; Fig. 2). For  $\delta^{15}\text{N}$ , the LMEM revealed significant differences  
345 between the Control and High treatment ( $12.4 \pm 0.6$  vs.  $10.6 \pm 1.0\text{‰}$ ;  $P < 0.001$ ), but not  
346 between the Control and the Low and Medium treatments ( $12.4 \pm 0.6$  vs.  $12.0 \pm 1.6$  and  $11.6$   
347  $\pm 1.6\text{‰}$  respectively;  $P = 1.0$  in all cases; Fig. 2). The 95% confidence limits of the estimates  
348 of isotopic niche size ( $\text{SEA}_b$ ) indicated that the niche of the *B. barbuis* in the low treatment  
349 was significantly larger than the Control, Medium and High treatments (Table 1; Fig. 2). The  
350 isotopic niche of the Control overlapped with that of the Low treatment by 76%, but did not  
351 overlap at all with the Medium and High treatments (Table 1; Fig. 2). In the Control,

352 macroinvertebrates were the principal contributor to *B. barbuis* diet, whereas in the Medium  
353 and High treatments, pellets contributed up to 48% of diet (Table 1). In the Low treatment,  
354 pellets only contributed 23% to estimated diet (Table 1).

355

#### 356 *Pond experiments*

357 Across the treatments, the mean starting lengths of the *B. barbuis* were 77.5 to 82.0 mm and *S.*  
358 *cephalus* 73.9 to 81.7 mm (Table 2). At the conclusion of the experiment, 97% of the fish  
359 present at the start of the experiment were recovered at the end (174 from 180 fish), with no  
360 more than one fish per replicate missing. The length range of the fish had increased to 113.7  
361 to 119.4 mm (*B. barbuis*) and 124.6 to 131.1 mm (*S. cephalus*). The generalized linear model  
362 testing differences in IL across the species and treatments was significant (Wald  $\chi^2 = 105.4$ ,  $P$   
363 = 0.02), with the effect of starting length being a significant covariate ( $P = 0.04$ ). Pairwise  
364 comparisons revealed, however, that there were no significant differences in growth rates  
365 across the species and their treatments ( $P = 0.09$  to 1.0; Fig. 3).

366

367 The LMEM revealed that the significant differences in the corrected  $\delta^{13}\text{C}$  data (Ccorr) were  
368 primarily between the pellet and no pellet treatments, including between allopatric *B. barbuis*  
369 (pellet:  $1.92 \pm 0.09$ ; no pellet:  $0.68 \pm 0.09$ ;  $P < 0.001$ ) and allopatric *S. cephalus* (pellet:  $1.84$   
370  $\pm 0.09$ ; no pellet:  $0.25 \pm 0.09$ ;  $P < 0.001$ ) (Fig. 4). The same differences were also apparent  
371 for TP, but with additional differences between the two fishes in the presence and absence of  
372 pellets ( $P < 0.02$  in all cases), where *B. barbuis* were at a higher TP than *S. cephalus* (Fig. 4).  
373 Isotopic niche estimates revealed that there was no overlap in the niches of the two fishes in  
374 allopatry or sympatry, or in the presence and absence of pellets, but the availability of pellets  
375 caused a substantial shift in the position of the isotopic niche of both fishes in both allopatry  
376 and sympatry (Fig. 4). This shift was caused by the presence of the pellets in fish diet; where



377 present, their contribution to fish diet was 43 and 58% (Table 3). In terms of isotopic niche  
378 size, however, there was considerable overlap in the 95% confidence limits of estimates of  
379  $SEA_b$  for the species in the presence/ absence of pellets in their allopatric and sympatric  
380 contexts, thus the pellets did not affect isotopic niche size (Table 4).

381

### 382 *Wild fishes*

383 A total of 31 *B. barbuis* were sampled from the River Teme in September 2015. Of these, 19  
384 were captured by electric fishing (mean length  $512.1 \pm 63.8$  mm) and 12 by angling (mean  
385 length  $616.8 \pm 72.7$  mm), with the differences in their lengths being significant (ANOVA:  
386  $F_{1,29} = 5.56$ ,  $P = 0.03$ ). Across this dataset, there was also a significant relationship between  
387 fish length and SI data ( $\delta^{13}C$ :  $R^2 = 0.42$ ,  $F_{1,29} = 20.61$ ,  $P < 0.001$ ;  $\delta^{15}N$ :  $R^2 = 0.32$ ,  $F_{1,29}$   
388  $= 13.50$ ,  $P < 0.001$ ). To remove this ontogenetic influence of length on the SI data, the six fish  
389 captured by electric fishing of  $< 400$  mm length were removed from the dataset, resulting in  
390 the relationships between fish length and SI data now being non-significant ( $\delta^{13}C$ :  $R^2 = 0.10$ ,  
391  $F_{1,23} = 2.30$ ,  $P = 0.13$ ;  $\delta^{15}N$ :  $R^2 = 0.09$ ,  $F_{1,23} = 2.18$ ,  $P = 0.15$ ). This also increased the mean  
392 length of the electric fished *B. barbuis* to  $585.8 \pm 55.9$  mm ( $n = 13$ ), with this not significantly  
393 different to the angler caught fish (ANOVA:  $F_{1,23} = 0.96$ ,  $P = 0.34$ ). In addition, 6 *S. cephalus*  
394 were sampled by angling (length range: 400 to 540 mm; mean length  $456.7 \pm 51.3$  mm), with  
395 none sampled by electric fishing. Regarding the age of the *B. barbuis*  $> 400$  mm, there was  
396 only one individual age at 8+ years, with the remainder all between 11+ and 18+ years. At  
397 these ages, their annual length increments were relatively low (mean last annual length  
398 increment:  $18.7 \pm 4.1$  mm), with the relationship between length increment and the SI data  
399 being non-significant ( $\delta^{13}C$ :  $R^2 = 0.04$ ,  $F_{1,23} = 0.67$ ,  $P = 0.42$ ;  $\delta^{15}N$ :  $R^2 = 0.08$ ,  $F_{1,23} = 1.56$ ,  $P$   
400  $= 0.23$ .

401

402 For the *B. barbuis* > 400 mm sampled by electric fishing, their isotopic niche was  
403 significantly larger than the angled fish (95% CL SEA<sub>b</sub>: 2.54 to 6.66 vs. 0.66 to 2.30‰; Fig.  
404 5). The angled sub-set of *B. barbuis* shared 83% of their isotopic space with those that were  
405 electric fished (Fig. 5). The angled *S. cephalus* had an isotopic niche in a similar position to  
406 the angled *B. barbuis* and they also had a similar niche size (95% CL SEA<sub>b</sub>: 0.63 to 4.28‰;  
407 Fig. 5). The estimated dietary contributions from the Bayesian mixing models suggested that  
408 the angled *B. barbuis* and *S. cephalus* had total contributions of pellets of 59 and 44%  
409 respectively, whereas this was reduced to 39% for the electric fished individuals of > 400 mm  
410 (Table 5a). At the individual level, estimated dietary proportions varied by sampling method,  
411 but with generally lower proportions of pellets in the diet of electric fished *B. barbuis* (range 9  
412 to 62%) than angled (range 40 to 71%) (Table 5b). The coefficient of variation was also  
413 higher for all food items for electric fished *B. barbuis*, but this was especially strong for  
414 pellets (electric fished: 0.45; angled: 0.17; Table 5b). The overall range of the contribution of  
415 pellets to *B. barbuis* diet, irrespective of sampling method, was 9 to 71% (Table 5b).

416

## 417 **Discussion**

418

419 The two experiments revealed that where fishmeal pellets were present as a food resource for  
420 *B. barbuis* and *S. cephalus*, these were generally consumed in sufficient proportions to alter  
421 the SI signatures of their tissues, as per the hypothesis, and resulted in major shifts in the  
422 position of their population isotopic niche. In wild *B. barbuis*, where fish were sampled by  
423 both angling and electric fishing, there was considerable individual variability in the  
424 contribution of pellets to diet, ranging between 9 and 71%; where only angled fish were  
425 considered then the range was 40 to 71%. High estimates of contributions of pellets to *S.*  
426 *cephalus* diet were also apparent, with these all captured by angling. The largest isotopic

427 niches were apparent in the ‘Low’ treatment of the mesocosm experiment and in the wild *B.*  
428 *barbus* captured by both angling and electric fishing. This was likely to be the result of the  
429 diets of the individual fish comprising of a greater variety of dietary items, in which MDN  
430 pellets were important items for only some individuals. Regarding somatic growth rates,  
431 whilst these were significantly higher in the ‘medium’ and ‘high’ treatments compared to the  
432 control and ‘low’ treatment in the mesocosm experiment, there were no significant  
433 differences in the growth rates of the fishes detected in the pond experiment, and there was  
434 no relationship between annual length increments and the SI data for the wild fishes. Thus,  
435 despite the pellets being consumed and assimilated into the fish tissues across the study  
436 approaches, it was only in very controlled conditions where feeding on pellets facilitated  
437 faster growth rates, and then only when they were available in relatively high quantities. This  
438 finding was generally contrary to the hypothesis.

439

440 Recent studies have suggested that where *B. barbus* populations are enhanced with hatchery  
441 reared individuals via stocking then there are strong patterns in isotopic niche partitioning  
442 between these fish and other wild fishes, including *S. cephalus* (Bašić & Britton, 2016). This  
443 partitioning is also evident between larger individuals, suggesting functional differences  
444 between the species result in these trophic differences (Bašić & Britton, 2015, 2016). This  
445 isotopic niche partitioning between *B. barbus* and *S. cephalus* was also apparent here, with  
446 the species having distinct niches in the presence and absence of pellets. Thus, even where  
447 the fishes feed on pellets in relatively high proportions, such as in the ‘pellet pond’ of the  
448 pond experiments, their functional differences were still sufficient to result in differences in  
449 the position of their isotopic niches. Reasons for these inter-specific isotopic niches  
450 differences might relate to differences in the proportions of macroinvertebrates consumed  
451 between the species and differences in the stable isotope ecology between *B. barbus* and *S.*

452 *cephalus*, for example through differences in their fractionation factors (Busst, Bašić &  
453 Britton, 2015; Busst & Britton, 2016). Irrespective, in this pond experiment, the growth rates  
454 and sizes of the isotopic niches of both fishes were not significantly different between their  
455 allopatric and sympatric contexts in both pellet presence and absence, suggesting that the  
456 fishes were accessing sufficient food resources to maintain their growth rates without having  
457 to further alter their diet.

458

459 It was apparent that all of the fish sampled by angling from the River Teme, both here and in  
460 Bašić *et al.* (2015), generally had diets comprising relatively high proportions of MDN (up to  
461 80% in Bašić *et al.* 2015), yet for *B. barbuis* sampled by electric fishing, there was much  
462 greater variability in this MDN contribution, with this independent of body size. This  
463 suggests that despite the attractiveness of fishmeal pellets to *B. barbuis* generally, resulting in  
464 some individuals developing trophic specialisations, other individuals primarily consumed  
465 other items, perhaps through avoiding consuming pellets due to previous angler capture  
466 experiences that lead to avoidance (Raaij, 1985; Askey *et al.*, 2006). This also emphasises the  
467 potential bias that can result from samples collected by angling alone, as individual  
468 variability in the behaviour of individuals can affect capture susceptibility (Klefoth *et al.*,  
469 2013).

470

471 It was apparent that the MDN from the pellets was being consumed directly by the fishes,  
472 with the stable isotope data of the macroinvertebrates and fish suggesting there was no  
473 indirect transfer via prey populations. This is in contrast to the transfer of MDN into  
474 freshwaters via migratory salmonid fishes, where the nutrients are more freely available and  
475 facilitate the increased production of benthic algae and macroinvertebrates (Schindler *et al.*,  
476 2003). This then enhances the food resources available for the larvae and juveniles of the

477 adult migrants, facilitating their feeding, growth and survival in the early life stages (Wipfli *et*  
478 *al.*, 2003). The MDN from salmonids can thus be traced through freshwater food webs,  
479 enabling assessment of the links between the aquatic and terrestrial food webs. For example,  
480 Tonra *et al.* (2015) reported on the removal of Elwha River dam in the USA, which resulted  
481 in migratory salmonids returning to the river within 12 months. Following reproduction and  
482 death of these fishes, their MDN could be traced through the macroinvertebrate community  
483 and then into a bird that preys upon these, the American dipper *Cinclus mexicanus*. Indeed,  
484 there are now numerous studies that have traced MDN into terrestrial food webs (e.g.  
485 McLoughlin *et al.*, 2016; Richardson *et al.*, 2016), with its influence even affecting the  
486 behaviour of terrestrial predator and scavenger species (Schindler *et al.*, 2013).

487

488 In contrast, the apparent direct transfer of MDN from fishmeal pellet to *B. barbuis* and *S.*  
489 *cephalus* in this study suggested that this nutrient subsidy might have only minor impacts on  
490 the non-fish communities. In the wild, the fish consuming these pellets tend to be large-  
491 bodied and thus are only likely to be predated upon by large piscivores, including otter *Lutra*  
492 *lutra*, although otters tend to prefer to consume high abundances of smaller bodied fishes  
493 (Britton *et al.*, 2006). Unlike salmonid fishes, *B. barbuis* and *S. cephalus* are relatively long-  
494 lived (> 15 years; Britton, 2007; Britton *et al.* 2013), reproducing annually following sexual  
495 maturity (Britton & Pegg, 2011), and thus there is no large post-spawning die-off.  
496 Consequently, they might be acting as MDN sinks, with low rates of nutrient transfer to  
497 higher trophic levels. However, determining the extent of MDN transfer to higher trophic  
498 levels requires further work. There might also be some alternative ecological benefits of this  
499 MDN subsidy. For example, in many European rivers, including the River Teme, *B. barbuis* is  
500 a large-bodied invasive fish that potentially impacts prey populations and competes with  
501 functional analogues (Antognazza *et al.*, 2016). Whilst recent studies suggest some trophic

502 (isotopic) partitioning between *B. barbus* and other fishes in riverine communities (Bašić &  
503 Britton, 2015, 2016), the high proportion of fishmeal pellets detected in the diet of wild  
504 fishes, both here and in Bašić *et al.* (2015), suggests this trophic subsidy could potentially  
505 lead to further partitioning between fish populations across the fish communities. This is also  
506 likely to reduce invasive *B. barbus* predation pressure on macroinvertebrate communities, as  
507 their dietary requirements are primarily met by the consumption of this angler subsidy.

508

509 These results add to an increasing literature base on the role of subsidies from fishery  
510 activities in the trophic ecology of freshwater communities. For example, Grey, Waldron &  
511 Hutchinson (2004) demonstrated that approximately 65% of *Daphnia* spp. and over 80% of  
512 roach *Rutilus rutilus* body carbon was ultimately derived from pellet material originating  
513 from an *in situ* fish farm in Esthwaite Water, England. These data suggest that the MDN were  
514 more freely available within the lake via the breakdown of the pellets, with a number of other  
515 studies also revealing their integration into the food web more generally (Fernandez-Jover *et*  
516 *al.*, 2011a,b; Demétrio *et al.*, 2012; Jackson *et al.*, 2013). Thus, further work is suggested in  
517 riverine systems where fishmeal pellets are used by anglers to identify whether there is  
518 greater transfer of MDN in the food web than suggested here.

519

520 In summary, across three spatial scales of increasing complexity, it was apparent that the  
521 release of fishmeal pellets into freshwaters as an allochthonous trophic subsidy based on  
522 MDN had a substantial influence on the isotopic niche (as a proxy of the trophic niche) of  
523 riverine fishes. Results from wild *B. barbus*, with some support from the experiments,  
524 indicated that individual isotopic niche specialisation resulting from this trophic subsidy was  
525 strongly apparent, with its development potentially associated with behavioural differences

526 between individual fish that leads to variability in their avoidance/ consumption of pellets and  
527 thus their likelihood of angler capture.

528

## 529 **Acknowledgments**

530

531 CGR was supported by a studentship funded by the Severn Rivers Trust and TB was  
532 supported by a studentship funded by the Environment Agency and Barbel Society. F.A-T  
533 holds a doctoral fellowship from the Spanish Ministry of Education (FPU13/00235). We  
534 specifically thank Brecht Morris, Alan Henshaw and Pete Reading for their assistance in  
535 obtaining samples from the River Teme. All work was completed under project licence and  
536 personal licences from the UK Home Office.

537

## 538 **References**

539

540 Antognazza C.M., Andreou D., Zaccara S. & Britton J.R. (2016) Loss of genetic integrity and  
541 biological invasions result from stocking and introductions of *Barbus barbus*: insights  
542 from rivers in England. *Ecology and Evolution*, **6**, 1280-1292.

543 Araújo M.S., Bolnick D.I. & Layman C.A. (2011) The ecological causes of individual  
544 specialisation. *Ecology Letters*, **14**, 948-958.

545 Arlinghaus R. & Niesar N. (2005) Nutrient digestibility of angling baits for carp, *Cyprinus*  
546 *carpio*, with implications for groundbait formulation and eutrophication control. *Fisheries*  
547 *Management and Ecology*, **12**, 91-97

548 Askey P.J., Richards S.A., Post J.R. & Parkinson E.A. (2006) Linking angling catch rates and  
549 fish learning under catch-and-release regulations. *North American Journal of Fisheries*  
550 *Management*, **26**, 1020-1029.

551 Bašić T. & Britton J.R. (2015) Utility of fish scales from stock assessment surveys in stable  
552 isotope analysis for initial assessments of trophic relationships in riverine fish  
553 communities. *Journal of Applied Ichthyology*, **31**, 296-300.

554 Bašić T., Britton J.R., Jackson M.C., Reading P. & Grey J. (2015) Angling baits and invasive  
555 crayfish as important trophic subsidies for a large cyprinid fish. *Aquatic Sciences*, **77**, 153-  
556 160.

557 Bašić T. & Britton J.R. (2016) Characterising the trophic niches of stocked and resident  
558 cyprinid fishes: consistency in partitioning over time, space and body sizes. *Ecology and  
559 Evolution* **6**, 5093-5104.

560 Britton J.R., Pegg J., Shepherd J.S. & Toms S. (2006) Revealing the prey items of the otter  
561 *Lutra lutra* in South West England using stomach contents analysis. *Folia Zoologica*, **55**,  
562 167-174

563 Britton J.R. (2007) Reference data for evaluating the growth of common riverine fishes in the  
564 UK. *Journal of Applied Ichthyology*, **23**, 555-560.

565 Britton J.R. & Pegg J. (2011) Ecology of European Barbel *Barbus barbus*: Implications for  
566 river, fishery, and conservation management. *Reviews in Fisheries Science*, **19**, 321-330.

567 Britton J.R., Davies G.D. and Pegg J. (2013) Spatial variation in the somatic growth rates of  
568 European barbel *Barbus barbus*: a UK perspective. *Ecology of Freshwater Fish*, **22**, 21-  
569 29.

570 Britton J.R. & Andreou D. (2016) Parasitism as a Driver of Trophic Niche Specialisation.  
571 *Trends in Parasitology*, **32**, 437-445.

572 Busst G., Bašić T. & Britton J.R. (2015) Stable isotope signatures and trophic-step  
573 fractionation factors of fish tissues collected as non-lethal surrogates of dorsal muscle.  
574 *Rapid Communications in Mass Spectrometry*, **29**, 1535-1544.

575 Busst G.M. & Britton J.R. (2016) High variability in stable isotope diet–tissue discrimination



576 factors of two omnivorous freshwater fishes in controlled ex situ conditions. *Journal of*  
577 *Experimental Biology*, **219**, 1060-1068.

578 Demétrio J.A., Gomes L.C., Latini J.D. & Agostinho A.A. (2012) Influence of net cage  
579 farming on the diet of associated wild fish in a Neotropical reservoir. *Aquaculture*, **330-**  
580 **333**, 172-178.

581 Dynamite Baits (2017) Marine Halibut Pellets.  
582 <http://www.dynamitebaits.com/products/p/marine-halibut-pellets>. Last accessed  
583 17/01/2017.

584 Fernandez-Jover D., Arechavala-Lopez P., Martinez-Rubio L., Tocher D.R., Bayle-Sempere  
585 J.T., Lopez-Jimenez J.A., Martinez-Lopez F.J. & Sanchez-Jerez P. (2011a) Monitoring the  
586 influence of marine aquaculture on wild fish communities: benefits and limitations of fatty  
587 acid profiles. *Aquaculture Environment Interactions*, **2**, 39-47.

588 Fernandez-Jover D., Martinez-Rubio L., Sanchez-Jerez P., Bayle-Sempere J.T., Lopez  
589 Jimenez J.A., Martínez Lopez F.J., Bjørn P.A., Uglem I. & Dempster T. (2011b) Waste  
590 feed from coastal fish farms: a trophic subsidy with compositional side-effects for wild  
591 gadoids. *Estuarine and Coastal Shelf Science*, **91**, 559-568.

592 Grey J., Waldron S. & Hutchinson R. (2004) The utility of carbon and nitrogen isotope  
593 analyses to trace contributions from fish farms to the receiving communities of freshwater  
594 lakes: a pilot study in Esthwaite Water, UK. *Hydrobiologia*, **524**, 253-262.

595 Grey J., Graham C.T., Britton J.R. & Harrod C. (2009) Stable isotope analysis of archived  
596 roach (*Rutilus rutilus*) scales for retrospective study of shallow lake responses to nutrient  
597 reduction. *Freshwater Biology*, **54**, 1663-1670.

598 Hobson K.A. & Clark R.G. (1992) Assessing avian diets using stable isotopes I: turnover of  
599 <sup>13</sup>C in tissues. *Condor*, **94**, 181-188.

600 Hutchinson J.J. & Trueman C.N. (2006) Stable isotope analyses of collagen in fish scales:  
601 limitations set by scale architecture. *Journal of Fish Biology*, **69**, 1874-1880.

602 Jackson A.L., Inger R., Parnell A.C. & Bearhop S. (2011) Comparing isotopic niche widths  
603 among and within communities: SIBER–Stable Isotope Bayesian Ellipses in R. *Journal of*  
604 *Animal Ecology*, **80**, 595-602.

605 Jackson M.C., Donohue I., Jackson A.L., Britton J.R., Harper D.M. & Grey J. (2012)  
606 Population-level metrics of trophic structure based on stable isotopes and their application  
607 to invasion ecology. *PLoS ONE* **7**:e31757.

608 Jackson M.C., Allen R., Pegg J. & Britton J.R. (2013) Do trophic subsidies affect the  
609 outcome of introductions of a non-native freshwater fish? *Freshwater Biology*, **58**, 2144-  
610 2153.

611 Jefferies R.L. (2000) Allochthonous inputs: integrating population changes and food-web  
612 dynamics. *Trends in Ecology and Evolution*, **15**, 19-22.

613 Jones R.I., Grey J., Sleep D. & Quarmby C. (1998) An assessment, using stable isotopes, of  
614 the importance of allochthonous organic carbon sources to the pelagic food web in Loch  
615 Ness. *Proceedings of the Royal Society of London: Biological Sciences*, **265**, 105-110.

616 Klefoth T., Pieterek T. and Arlinghaus R. (2013) Impacts of domestication on angling  
617 vulnerability of common carp, *Cyprinus carpio*: the role of learning, foraging behaviour  
618 and food preferences. *Fisheries Management and Ecology*, **20**, 174-186.

619 Marcarelli A.M., Baxter C.V., Mineau M.M. & Hall R.O. (2011) Quantity and quality:  
620 unifying food web and ecosystem perspectives on the role of resource subsidies in  
621 freshwaters. *Ecology*, **92**, 1215-1225.

622 Marczak L.B., Thompson R.M. & Richardson, J.S. (2007) Meta-analysis: trophic level,  
623 habitat, and productivity shape the food web effects of resource subsidies. *Ecology*, **88**,  
624 140-148.

625 McLoughlin P.D., Lysak K., Debeffe L., Perry T. & Hobson K.A. (2016) Density-dependent  
626 resource selection by a terrestrial herbivore in response to sea-to-land nutrient transfer by  
627 seals. *Ecology*, **97**, 1929-1937.

628 Naylor R.L., Goldburg R.J., Primavera J.H., Kautsky N., Beveridge M.C.M., Clay J., Folke  
629 C., Lubchenco J., Mooney H. & Troell M. (2000) Effect of aquaculture on world fish  
630 supplies. *Nature*, **405**, 1017-1024.

631 Olsson K., Stenroth P., Nyström P. & Graneli W. (2009) Invasions and niche width: Does  
632 niche width of an introduced crayfish differ from a native crayfish? *Freshwater Biology*,  
633 **54**, 1731–1740.

634 Parnell A.C., Inger R., Bearhop S. & Jackson A.L. (2010) Source partitioning using stable  
635 isotopes: Coping with too much variation. *PLoS ONE*, **5**, e9672.

636 Polis G.A. & Hurd S.D. (1995) Extraordinarily high spider densities on islands: flow of  
637 energy from the marine to terrestrial food webs and the absence of predation. *Proceedings*  
638 *of the National Academy of Sciences USA*, **92**, 4382-4386.

639 Post D.M. (2002) Using stable isotopes to estimate trophic position: Models, methods, and  
640 assumptions. *Ecology*, **83**, 703-718.

641 Post D.M., Layman C.A., Arrington D.A., Takimoto G., Quattrochi J. & Montana C.G.  
642 (2007) Getting to the fat of the matter: models, methods and assumptions for dealing with  
643 lipids in stable isotope analyses. *Oecologia*, **152**, 179-189.

644 R Development Core Team (2013) R: A language and environment for statistical computing.

645 Raat A.J.P. (1985) Analysis of angling vulnerability of common carp, *Cyprinus carpio* L., in  
646 catch-and-release angling in ponds. *Aquaculture Research*, **16**, 171-187.

647 Richardson D.P., Kohler A.E., Hailemichael M. & Finney B.P. (2016) The fate of marine-  
648 derived nutrients: tracing  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  through oligotrophic freshwater and linked

649 riparian ecosystems following salmon carcass analog additions. *Canadian Journal of*  
650 *Fisheries and Aquatic Sciences*, **73**, 1-15.

651 Sato T. & Watanabe K. (2013) Do stage-specific functional responses of consumers dampen  
652 the effects of subsidies on trophic cascades in streams? *Journal of Animal Ecology*, **83**,  
653 907-915.

654 Schindler D.E., Scheuerell M.D., Moore J.W., Gende S.M., Francis T.B. & Palen W.J. (2003)  
655 Pacific salmon and the ecology of coastal ecosystems. *Frontiers in Ecology and the*  
656 *Environment*, **1**, 31-37.

657 Schindler D.E., Leavitt P.R., Brock C.S., Johnson S.P. & Quay P.D. (2005) Marine-derived  
658 nutrients, commercial fisheries and production of salmon and lake algae in Alaska.  
659 *Ecology*, **86**, 3225-3231.

660 Schindler D.E., Armstrong J.B., Bentley K.T., Jankowski K., Lisi P.J. & Payne L.X. (2013)  
661 Riding the crimson tide: mobile terrestrial consumers track phenological variation in  
662 spawning of an anadromous fish. *Biology Letters*, **9**, p.20130048.

663 Stock B.C. & Semmens B.X. (2013). MixSIAR GUI User Manual. Version 3.1.  
664 <https://github.com/brianstock/MixSIAR/>. doi:10.5281/zenodo.47719. Last accessed  
665 17/09/2016.

666 Syrjanen J., Korsu K., Louhi P., Paavola R. & Muotka T. (2011) Stream salmonids as  
667 opportunistic foragers: the importance of terrestrial invertebrates along a stream-size  
668 gradient. *Canadian Journal of Fisheries and Aquatic Sciences*, **68**, 2146-2156.

669 Thomas S.M. & Crowther T.W. (2015) Predicting rates of isotopic turnover across the animal  
670 kingdom: a synthesis of existing data. *Journal of Animal Ecology*, **84**, 861-870.

671 Tonra C.M., Sager-Fradkin K., Morley S.A., Duda J.J. & Marra P.P. (2015) The rapid return  
672 of marine-derived nutrients to a freshwater food web following dam removal. *Biological*  
673 *Conservation*, **192**, 130-134.

- 674 Tran T.N.Q., Jackson M.C., Sheath D., Verreycken H. & Britton J.R. (2015) Patterns of  
675 trophic niche divergence between invasive and native fishes in wild communities are  
676 predictable from mesocosm studies. *Journal of Animal Ecology*, **84**, 1071–1080.
- 677 Wipfli M.S., Hudson J.P., Caouette J.P. & Chaloner D.T. (2003) Marine subsidies in  
678 freshwater ecosystems: salmon carcasses increase the growth rates of stream-resident  
679 salmonids. *Transactions of the American Fisheries Society*, **132**, 371-381.
- 680 Zhang Y., Negishi J.N., Richardson J.S. & Kolodziejczyk R. (2003) Impacts of marine-  
681 derived nutrients on stream ecosystem functioning. *Proceedings of the Royal Society of  
682 London B: Biological Sciences*, **270**, 2117-2123.

Table 1. Mean lengths and weights, isotopic niche size (as 95% CL of standard ellipse area, SEA<sub>b</sub>) of *Barbus barbuis* per treatment and the extent of their overlap between treatments, and the estimated contributions of putative foods to their diet (0 – 1 scale), as predicted in MixSIAR ( $\pm 95\%$  CL). Sample sizes were n = 15 per treatment.

Treatment	Mean length (mm)		Mean weight (g)		SEA <sub>b</sub> (‰)	Overlap in isotopic niche with Control (%)	Estimated contribution to diet (%)	
	Start	End	Start	End			Macroinvertebrate	Pellet
Control	106.5 ± 8.5	108.2 ± 8.3	9.9 ± 1.8	11.2 ± 2.2	0.06 – 0.21	n /a	0.97 ± 0.02	0.03 ± 0.02
Low	103.8 ± 5.9	113.3 ± 6.6	10.2 ± 1.2	14.7 ± 2.5	0.39 – 1.31	76	0.77 ± 0.02	0.23 ± 0.02
Medium	105 ± 3.9	127.3 ± 3.9	12.3 ± 1.0	22.9 ± 2.5	0.10 – 0.33	0	0.52 ± 0.02	0.48 ± 0.02
High	106.6 ± 4.1	132.7 ± 6.6	11.6 ± 0.9	24.3 ± 3.4	0.08 – 0.28	0	0.54 ± 0.02	0.47 ± 0.02

Table 2. Number of fish per species and treatment analysed for stable isotope analysis from the pond enclosure experiment, their start and end mean lengths ( $\pm$  95% CL), and mean stable isotope values ( $\pm$  95% CL).

Treatment	Species	n	Mean starting length (mm)	Mean end length (mm)	Mean $\delta^{13}\text{C}$ (‰)	Mean $\delta^{15}\text{N}$ (‰)
Allopatry/pellets	<i>B. barbuis</i>	18	80.1 $\pm$ 0.3	117.83 $\pm$ 1.99	-24.70 $\pm$ 0.21	9.39 $\pm$ 0.10
Allopatry/pellets	<i>S. cephalus</i>	18	81.7 $\pm$ 0.4	131.06 $\pm$ 1.38	-25.10 $\pm$ 0.23	8.44 $\pm$ 0.04
Allopatry/no pellets	<i>B. barbuis</i>	18	77.6 $\pm$ 0.2	113.67 $\pm$ 1.32	-28.20 $\pm$ 0.20	11.18 $\pm$ 0.05
Allopatry/no pellets	<i>S. cephalus</i>	17	73.9 $\pm$ 0.3	124.59 $\pm$ 1.69	-30.31 $\pm$ 0.19	10.72 $\pm$ 0.05
Sympatry/pellets	<i>B. barbuis</i>	15	82.0 $\pm$ 0.4	119.4 $\pm$ 1.84	-25.45 $\pm$ 0.18	9.25 $\pm$ 0.09
Sympatry/pellets	<i>S. cephalus</i>	15	76.3 $\pm$ 0.4	125.27 $\pm$ 1.69	-24.94 $\pm$ 0.20	8.34 $\pm$ 0.04
Sympatry/no pellets	<i>B. barbuis</i>	15	77.5 $\pm$ 0.3	118.94 $\pm$ 1.91	-29.05 $\pm$ 0.11	10.79 $\pm$ 0.05
Sympatry/no pellets	<i>S. cephalus</i>	15	76.1 $\pm$ 0.4	126.73 $\pm$ 1.64	-30.67 $\pm$ 0.14	10.81 $\pm$ 0.03

Table 3. Estimated contributions (0 – 1) of each putative food item to fish diet in the ‘pellet’ treatments of the pond enclosure experiment. Values represent mean estimated dietary proportions ( $\pm$  95% CL) from MixSIAR.

	Corixidae	Odonata	2mm pellet	3mm pellet	Total pellet*
Allopatric <i>B. barbuis</i> (n=18)	0.34 $\pm$ 0.11	0.21 $\pm$ 0.13	0.27 $\pm$ 0.06	0.18 $\pm$ 0.06	0.45
Allopatric <i>S. cephalus</i> (n=15)	0.26 $\pm$ 0.04	0.16 $\pm$ 0.05	0.33 $\pm$ 0.04	0.25 $\pm$ 0.04	0.58
Sympatric <i>B. barbuis</i> (n=18)	0.32 $\pm$ 0.11	0.22 $\pm$ 0.12	0.25 $\pm$ 0.06	0.22 $\pm$ 0.07	0.47
Sympatric <i>S. cephalus</i> (n=15)	0.25 $\pm$ 0.09	0.15 $\pm$ 0.10	0.33 $\pm$ 0.09	0.27 $\pm$ 0.11	0.60

\* derived from additional of the modal estimations of the 2mm and 3mm pellet and so no estimate of error around the values are provided.



Table 4. Isotopic niche size, as 95% CL of SEA<sub>b</sub> (‰) for *Barbus barbuis* and *Squalius cephalus* in the different treatments of the pond enclosure experiment, and as calculated from corrected stable isotope data. Sample sizes were as per Table 3.

	n	No fishmeal pellet	Fishmeal pellet
Allopatric <i>B. barbuis</i>	18	0.02 – 0.05	0.03 – 0.09
Sympatric <i>B. barbuis</i>	18	0.01 – 0.03	0.02 – 0.04
Allopatric <i>S. cephalus</i>	15	0.02 – 0.05	0.02 – 0.05
Sympatric <i>S. cephalus</i>	15	0.01 – 0.02	0.01 – 0.04

Table 5. (a) Mean contributions to fish diet of putative food resources (0 – 1 scale;  $\pm$  95% CL) of *Barbus barbuis* and *Squalius cephalus* in the River Teme by sampling method, estimated by MixSIAR; (b) minimum, maximum, mean ( $\pm$  95% CL) and coefficient of variation (CV) of estimates of contributions to individual *B. barbuis* diet (0 – 1) of the putative foods per sampling method (EF: electric fishing; A: angling), estimated by SOLOSIAR, where mean pellet data represents the sum of mean Pellet 1 and mean Pellet 2 per individual fish. Only *B. barbuis* of > 400 mm length were used in analyses.

(a)

Species	n	Arthropoda	'Small fishes'	Pellet 1	Pellet 2	Total pellet*
Electric fished <i>B. barbuis</i>	13	0.39 $\pm$ 0.10	0.26 $\pm$ 0.09	0.10 $\pm$ 0.04	0.26 $\pm$ 0.04	0.36
Angled <i>B. barbuis</i>	12	0.22 $\pm$ 0.07	0.20 $\pm$ 0.06	0.11 $\pm$ 0.03	0.48 $\pm$ 0.04	0.59
Angled <i>S. cephalus</i>	6	0.23 $\pm$ 0.11	0.24 $\pm$ 0.10	0.15 $\pm$ 0.06	0.39 $\pm$ 0.08	0.54

\* derived from additional of the modal estimations of the 2mm and 3mm pellet and so no estimate of error around the values are provided.

(b)

Dietary item	Minimum		Maximum		Mean		CV	
	EF	A	EF	A	EF	A	EF	A
Arthropod	0.07	0.13	0.45	0.30	0.19 $\pm$ 0.09	0.18 $\pm$ 0.05	0.82	0.68
Small fish	0.18	0.16	0.50	0.43	0.23 $\pm$ 0.10	0.24 $\pm$ 0.05	0.81	0.69
Pellet	0.09	0.40	0.62	0.71	0.38 $\pm$ 0.09	0.59 $\pm$ 0.06	0.45	0.17

## Figure captions

Figure 1. Somatic growth rates, as specific growth rate (A) and incremental length (B) per treatment for *Barbus barbuis* in the mesocosm experiment. Values represent estimated marginal means from the generalized linear models and \* indicates the difference in growth rate is significant at  $P < 0.001$ ) between the treatment and the control according to linearly independent pairwise comparisons. Error bars represent 95% confidence limits.

Figure 2. Stable isotope bi-plot of *Barbus barbuis* in the 250 L mesocosms and their isotopic niche (as standard ellipse area,  $SEA_c$ ), where clear triangles are the control fish and solid black line is their isotopic niche, filled triangles are the low treatment fish and the dashed black line is their isotopic niche, clear circles are the medium treatment fish and the solid light grey line is their isotopic niche, and grey circles are the high treatment fish and the dark grey line is their isotopic. × represent Chironomid larvae and + represent the fishmeal pellets fed daily.

Figure 3. Somatic growth rates, as incremental length, of *Barbus barbuis* (filled circles) and *Squalius cephalus* (clear circles) per treatment in the pond enclosure experiment. BAP: allopatric *B. barbuis* with pellets; BAN: allopatric *B. barbuis*, no pellets; BSP: sympatric *B. barbuis* with pellets; BSN: sympatric *B. barbuis*, no pellets; CAP: allopatric *S. cephalus* with pellets; CAN: allopatric *S. cephalus*, no pellets; CSP: sympatric *S. cephalus* with pellets; CSN: sympatric *S. cephalus*, no pellets. Error bars represent 95% confidence limits.

Figure 4. Stable isotope biplots (of corrected stable isotope data to trophic position and corrected carbon,  $C_{corr}$ ) showing individual data points (as symbols) and the isotopic niche (as standard ellipse area,  $SEA_c$ ) for (A) allopatric *Squalius cephalus* in the no pellet (clear

circle, solid black line) and pellet treatment (filled circle, dashed black line); (B) allopatric *Barbus barbatus* in the no pellet (clear square, solid grey line) and pellet treatment (filled square, dashed grey line); and (C) sympatric *S. cephalus* in the no pellet (clear circle, solid black line) and pellet treatment (filled circle, dashed black line), and sympatric *B. barbatus* in the no pellet (clear square, solid grey line) and pellet treatment (filled square, dashed grey line).

Figure 5. Stable isotope bi-plot of the lower River Teme, showing individual data points and isotopic niches (as standard ellipse areas). *Barbus barbatus* (electric fishing; length range 401 to 770 mm; n = 13): data points: black circles, solid black line: isotopic niche; *Barbus barbatus* (angling, length range 520 to 721 mm; n = 12): data points: clear circles, dashed black line: isotopic niche; *Squalius cephalus* (angling, length range 400 to 540 mm; n = 6): data points: clear squares, solid grey line: isotopic niche, Grey circles are combined data for 'small fishes' (*Cottus gobio*, *Barbatula barbatula*, *Phoxinus phoxinus*); + fishmeal pellet 1; × fishmeal pellet 2; black triangle: Arthropoda.




Figure 1.




Figure 2.




Figure 3.




Figure 4.






Figure 5.